

PS2.290

Doctor, want Meperidina please

Pilar Valverde Vallejo(1), J Palacios Castillo(2), D Ruiz Rueda(3)

(1) Hospital Regional de Málaga, Servicio de Urgencias, Málaga, Spain

(2) Hospital Regional de Málaga, Hospital Civil Servicio de Urgencias, Málaga, Spain

(3) Hospital Parque San Antonio, Servicio de Urgencias y Cuidados Críticos, Spain

Corresponding author: Dr Pilar Valverde Vallejo, Hospital Regional de Malaga, Urgencias, (Er), Malaga, Spain. E-mail: pilarvalverde79@gmail.com

Background and Purpose: Addiction to opioids is defined as a pattern of compulsive behavior that leads the individual who suffers a continuous search of the opioid and its use for other non-analgesic effects. Accordingly, some authors have defined other terms, the "pseudoaddiction" which is observed in patients with cancer or chronic pain, and in his search for opiates is due to a psychological dependence, but with analgesic purposes.

Addressing opioid dependence disorder in our hiperconsultant specifically in a patient with a sore arm and catastrophic present resolution after withdrawal

Method: Set the treatment of opioid dependence as a method of detoxification MEPERIDINE dependent patients. Detoxification goals are:

- 1) To release the body of the physical dependency associated with chronic consumption of narcotics.
- 2) Reduce or eliminate the pain and discomfort that may occur during withdrawal.
- 3) Provide safe and humane treatment to help the individual overcome the initial hurdle to stop using narcotics
- 4) Provide safe treatment that leads to a broader commitment to make appropriate treatment and these other treatment modalities leads.
- 5) Treat any medical problem that can be detected or perform the most appropriate referrals.
- 6) Start the process of educating the patient detoxification is successful when the patient is safe and with minimal discomfort.

Conclusions:

- The opioid dependence disorder is one of the most difficult to address in our area, because often seen in persons linked to the area onf health.
- While abstinence is the therapeutic goal of any dependence disorder, to achieve this there must be a prior strengthening the doctor-patient relationship and psychological readiness for change that the individual will perform.
- The treatment of opioid dependence involves not only physical detoxification, but also the psychological addiction, whether the therapeutic community or long-term outpatient treatment.

